

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

WEST Search History*Search Page 1*

DATE: Saturday, June 28, 2003

<u>Set Name</u> side by side	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u> result set
<i>DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ</i>			
L3	L2 and @ay<2001	42	L3
L2	L1 same activity	64	L2
L1	retro inverso peptide	155	L1

END OF SEARCH HISTORY

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1805JXB

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1		Web Page URLs for STN Seminar Schedule - N. America
NEWS 2		"Ask CAS" for self-help around the clock
NEWS 3	Jun 03	New e-mail delivery for search results now available
NEWS 4	Aug 08	PHARMAMarketLetter(PHARMAML) - new on STN
NEWS 5	Aug 19	Aquatic Toxicity Information Retrieval (AQUIRE) now available on STN
NEWS 6	Aug 26	Sequence searching in REGISTRY enhanced
NEWS 7	Sep 03	JAPIO has been reloaded and enhanced
NEWS 8	Sep 16	Experimental properties added to the REGISTRY file
NEWS 9	Sep 16	CA Section Thesaurus available in CAPLUS and CA
NEWS 10	Oct 01	CASREACT Enriched with Reactions from 1907 to 1985
NEWS 11	Oct 24	BEILSTEIN adds new search fields
NEWS 12	Oct 24	Nutraceuticals International (NUTRACEUT) now available on STN
NEWS 13	Nov 18	DKILIT has been renamed APOLLIT
NEWS 14	Nov 25	More calculated properties added to REGISTRY
NEWS 15	Dec 04	CSA files on STN
NEWS 16	Dec 17	PCTFULL now covers WP/PCT Applications from 1978 to date
NEWS 17	Dec 17	TOXCENTER enhanced with additional content
NEWS 18	Dec 17	Adis Clinical Trials Insight now available on STN
NEWS 19	Jan 29	Simultaneous left and right truncation added to COMPENDEX, ENERGY, INSPEC
NEWS 20	Feb 13	CANCERLIT is no longer being updated
NEWS 21	Feb 24	METADEx enhancements
NEWS 22	Feb 24	PCTGEN now available on STN
NEWS 23	Feb 24	TEMA now available on STN
NEWS 24	Feb 26	NTIS now allows simultaneous left and right truncation
NEWS 25	Feb 26	PCTFULL now contains images
NEWS 26	Mar 04	SDI PACKAGE for monthly delivery of multifile SDI results
NEWS 27	Mar 20	EVENTLINE will be removed from STN
NEWS 28	Mar 24	PATDPAFULL now available on STN
NEWS 29	Mar 24	Additional information for trade-named substances without structures available in REGISTRY
NEWS 30	Apr 11	Display formats in DGENE enhanced
NEWS 31	Apr 14	MEDLINE Reload
NEWS 32	Apr 17	Polymer searching in REGISTRY enhanced
NEWS 33	Jun 13	Indexing from 1947 to 1956 added to records in CA/CAPLUS
NEWS 34	Apr 21	New current-awareness alert (SDI) frequency in WPIDS/WPINDEX/WPIX
NEWS 35	Apr 28	RDISCLOSURE now available on STN
NEWS 36	May 05	Pharmacokinetic information and systematic chemical names added to PHAR
NEWS 37	May 15	MEDLINE file segment of TOXCENTER reloaded
NEWS 38	May 15	Supporter information for ENCOMPPAT and ENCOMPLIT updated
NEWS 39	May 16	CHEMREACT will be removed from STN
NEWS 40	May 19	Simultaneous left and right truncation added to WSCA
NEWS 41	May 19	RAPRA enhanced with new search field, simultaneous left and right truncation
NEWS 42	Jun 06	Simultaneous left and right truncation added to CBNB
NEWS 43	Jun 06	PASCAL enhanced with additional data
NEWS 44	Jun 20	2003 edition of the FSTA Thesaurus is now available
NEWS 45	Jun 25	HSDB has been reloaded

NEWS EXPRESS April 4 CURRENT WINDOWS VERSION IS V6.01a, CURRENT
 MACINTOSH VERSION IS V6.0b(ENG) AND V6.0Jb(JP),
 AND CURRENT DISCOVER FILE IS DATED 01 APRIL 2003
 NEWS HOURS STN Operating Hours Plus Help Desk Availability
 NEWS INTER General Internet Information
 NEWS LOGIN Welcome Banner and News Items
 NEWS PHONE Direct Dial and Telecommunication Network Access to STN
 NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
 specific topic.

All use of STN is subject to the provisions of the STN Customer
 agreement. Please note that this agreement limits use to scientific
 research. Use for software development or design or implementation
 of commercial gateways or other similar uses is prohibited and may
 result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 10:23:27 ON 28 JUN 2003

=> file .pub

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 10:23:59 ON 28 JUN 2003

FILE 'BIOSIS' ENTERED AT 10:23:59 ON 28 JUN 2003

COPYRIGHT (C) 2003 BIOLOGICAL ABSTRACTS INC. (R)

=> s retro inverso peptide

L1 96 RETRO INVERSO PEPTIDE

=> s l1 and review/dt

L2 0 L1 AND REVIEW/DT

=> s l1 and py<2001

L3 79 L1 AND PY<2001

=> duplicate remove l3

DUPLICATE PREFERENCE IS 'MEDLINE, BIOSIS'

KEEP DUPLICATES FROM MORE THAN ONE FILE? Y/(N):n

PROCESSING COMPLETED FOR L3

L4 52 DUPLICATE REMOVE L3 (27 DUPLICATES REMOVED)

=> d 1-10 bib ab

L4 ANSWER 1 OF 52 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

AN 2001:230910 BIOSIS

DN PREV200100230910

TI Probing SAR of FLRF-NH2 with its N- and C-terminally modified analogs and
retro-inverso peptides.

AU Kubiak, Teresa M. (1); Larsen, Martha J. (1); Dutton, Fred E. (1);
 Friedman, Alan R. (1)

CS (1) Animal Health Discovery Research, Pharmacia and Upjohn, Kalamazoo, MI,
 49001 USA

SO Fields, Gregg B.; Tam, James P.; Barany, George. (2000) pp. 762-763.
 Peptides for the new millennium. print.

Publisher: Kluwer Academic Publishers 3300 AA, Dordrecht, Netherlands.

Meeting Info.: 16th American Peptide Symposium Minneapolis, MI, USA June
 26-July 01, 1999

ISBN: 0-7923-6445-7 (cloth).
DT Book; Conference
LA English
SL English

L4 ANSWER 2 OF 52 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
AN 2000:417340 BIOSIS
DN PREV200000417340
TI Solid-phase synthesis of partially-modified retro and retro-inverso
psi(NHCH(CF₃))-peptides.
AU Volonterio, Alessandro (1); Bravo, Pierfrancesco; Moussier, Nathalie;
Zanda, Matteo (1)
CS (1) Dipartimento di Chimica del Politecnico, Via Mancinelli 7, I-20131,
Milano Italy
SO Tetrahedron Letters, (12 August, 2000) Vol. 41, No. 33, pp.
6517-6521. print.
ISSN: 0040-4039.

DT Article
LA English
SL English

AB The solid-phase synthesis of a novel class of retro and **retro-inverso peptides** featuring a psi(NHCH(CF₃)) surrogate of the classical (NH-CO) retro-peptide bond has been accomplished. Wang resin bound alpha-amino esters 2 were engaged in Michael-type N-additions with 3-(E-enoyl)-1,3-oxazolidin-2-one 3, which took place very effectively. Highly chemoselective exocyclic oxazolidinone cleavage, followed by parallel couplings of the resulting polymer bound pseudo-peptides 6 with further alpha-amino esters, and final release from the resins 7 delivered a library of nine psi(NHCH(CF₃)) retro and retro-inverso pseudo-tripeptides 8 with purity ranging from 75 to > 95%.

L4 ANSWER 3 OF 52 MEDLINE DUPLICATE 1
AN 2000483575 MEDLINE
DN 20457139 PubMed ID: 11000007

TI Design and solution structure of functional peptide mimetics of nerve growth factor.
AU Beglova N; Maliartchouk S; Ekiel I; Zaccaro M C; Saragovi H U; Gehring K
CS Department of Biochemistry and Montreal Joint Centre for Structural Biology, McGill University, 3655 Promenade Sir William Osler, Montreal, Quebec H3G 1Y6, Canada.
SO JOURNAL OF MEDICINAL CHEMISTRY, (2000 Sep 21) 43 (19) 3530-40.
Journal code: 9716531. ISSN: 0022-2623.

CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English

FS Priority Journals
EM 200010

ED Entered STN: 20001019
Last Updated on STN: 20001019
Entered Medline: 20001012

AB The C-D loop in nerve growth factor (NGF) is involved in binding to the NGF receptor, TrkA. It is flexible and adopts several different types conformations in different NGF crystal forms. We have previously shown that a small cyclic peptide derived from the C-D loop of NGF binds to the TrkA receptor by mimicking the structure of this loop. To understand structure-function relationships in NGF C-D loop mimetics, we have produced a series of peptides predicted to form different types of beta-turns. The peptides were tested for their ability to promote cell survival in serum-free medium and to induce TrkA tyrosine phosphorylation. NMR structural studies were used to determine the backbone conformation and the spatial orientation of side chains involved in binding to the TrkA receptor. Peptides that form type I or type gammaL-alphaR beta-turns were the most active. The variety of active loop conformations suggests that the mimetics (and NGF) accommodate the binding site on TrkA by an 'induced

fit' mechanism. In agreement with this hypothesis, NMR relaxation measurements detected both fast and slow motion in the peptides. We also characterized a **retro-inverso peptide** derived from the NGF C-D loop. This D-amino acid cyclic peptide did not adopt a conformation homologous to the NGF C-D loop and was inactive. This may be representative of difficulties in producing structural and functional mimetics by retro-inverso schemes.

L4 ANSWER 4 OF 52 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
AN 2001:78061 BIOSIS
DN PREV200100078061
TI A stable prosaposin **retro-inverso peptide**
exacerbates ischemia-induced behavioral deficits in rabbits: comparison
with the neuroprotective neurosteroid dehydroepiandrosterone sulfate.
AU Chapman, D. F. (1); Araujo, D. M.; Zivin, J. A.; Lapchak, P. A.
CS (1) UCSD, La Jolla, CA USA
SO Society for Neuroscience Abstracts, (2000) Vol. 26, No. 1-2, pp. Abstract
No.-287.5. print.
Meeting Info.: 30th Annual Meeting of the Society of Neuroscience New
Orleans, LA, USA November 04-09, 2000 Society for Neuroscience
. ISSN: 0190-5295.
DT Conference
LA English
SL English
AB Evidence suggests that prosaposin and neurosteroids can rescue neurons
from ischemic damage and excitotoxicity. Because of their potential
neuroprotective properties, both RIP and DHEAS may be useful in treating
ischemic stroke. We examined the behavioral effects of a stable 11-mer RIP
also known as Prosaptide (all D amino acids: LLEETANNDDL) and DHEAS in
rabbits exposed to reversible spinal cord ischemia produced by temporary
occlusion of the infrarenal aorta; RIP (1 mg/kg) or DHEAS (50 mg/kg) were
administered IV 5 minutes following various durations of aortic occlusion
ranging from 15 to 60 min, which allows for the calculation of the
duration (min) associated with a 50% probability of permanent paraplegia
(P50) for each experimental group. A drug was considered to be
neuroprotective only if it prolonged the P50 compared to the
vehicle-treated control group, which was approximately 25-28 min.
Treatment with RIP significantly ($p < 0.05$) decreased the P50 to 20 min (20%
reduction), whereas DHEAS significantly ($p < 0.05$) prolonged the P50 to 38
min (35% increase). The prominent neuroprotective effects that were
observed with DHEAS included increased mobility, tactile sensation and
hind limb use. In contrast, RIP exacerbated ischemia-induced behavioral
deficits and increased paraplegia. Overall, our study shows that although
neurotrophic-like properties have been documented for both RIP and DHEAS,
only the latter promotes recovery of spinal cord neuron function following
ischemia, suggesting that it may have therapeutic benefits for the
treatment of ischemic stroke.

L4 ANSWER 5 OF 52 MEDLINE DUPLICATE 2
AN 1999121111 MEDLINE
DN 99121111 PubMed ID: 9920919
TI Solution structure of a **retro-inverso peptide**
analogue mimicking the foot-and-mouth disease virus major antigenic site.
Structural basis for its antigenic cross-reactivity with the parent
peptide.
AU Petit M C; Benkirane N; Guichard G; Du A P; Marraud M; Cung M T; Briand J
P; Muller S
CS Laboratoire de Chimie-Physique Macromoleculaire, UMR 7568 CNRS,
ENSIC-INPL, 54000 Nancy, France.
SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1999 Feb 5) 274 (6) 3686-92.
Journal code: 2985121R. ISSN: 0021-9258.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English

FS Priority Journals
 OS PDB-1BCV; PDB-1BFW
 EM 199902
 ED Entered STN: 19990316
 Last Updated on STN: 20000303
 Entered Medline: 19990226
 AB The antigenic activity of a 19-mer peptide corresponding to the major antigenic region of foot-and-mouth disease virus and its retro-enantiomeric analogue was found to be completely abolished when they were tested in a biosensor system in trifluoroethanol. This suggests that the folding pattern, which is alpha-helix in trifluoroethanol (confirmed by CD measurement), does not correspond to the biologically relevant conformation(s) recognized by antibodies. The NMR structures of both peptides were thus determined in aqueous solution. These studies showed that the two peptides exhibit similar folding features, particularly in their C termini. This may explain in part the cross-reactive properties of the two peptides in aqueous solution. However, the retro-inverso analogue appears to be more rigid than the parent peptide and contains five atypical beta-turns. This feature may explain why retro-inverso foot-and-mouth disease virus peptides are often better recognized than the parent peptide by anti-virion antibodies.

L4 ANSWER 6 OF 52 MEDLINE DUPLICATE 3
 AN 1999388009 MEDLINE
 DN 99388009 PubMed ID: 10458771
 TI Inhibition of experimental autoimmune encephalomyelitis in SJL mice by oral administration of retro-inverso derivative of encephalitogenic epitope P87-99.
 AU Marino M; Ippolito A; Fassina G
 CS Biopharmaceuticals, TECNOGEN S.C.p.A., Science Park, Piana di Monte Verna, Italy.
 SO EUROPEAN JOURNAL OF IMMUNOLOGY, (1999 Aug) 29 (8) 2560-6.
 Journal code: 1273201. ISSN: 0014-2980.
 CY GERMANY: Germany, Federal Republic of
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199909
 ED Entered STN: 19990925
 Last Updated on STN: 20000303
 Entered Medline: 19990914
 AB Retro-inverso modification of peptides preserves parent peptide overall topology and provides at the same time stability to proteolysis, leading to derivatives with prolonged half-life in vitro and in vivo. In this study the encephalitogenic epitope P87 - 99 of myelin basic protein has been prepared in the retro-inverso form to examine its biological activity in a murine model of multiple sclerosis. Experiments of in vivo T cell tolerance induction in SJL mice revealed that the **retro-inverso peptide** was able to induce a selective T cell hyporesponsiveness, as measured by a reduction in the proliferative response of lymphnode T cells after antigen challenge. Oral administration of **retro-inverso peptide** decreased the disease severity significantly and delayed considerably the disease onset in treated mice. Enhancement of resistance to proteolysis by retro-inverso modification of encephalitogenic epitopes may increase the therapeutic value of oral tolerance induction in the treatment of multiple sclerosis and other Th1-associated inflammatory disorders.

L4 ANSWER 7 OF 52 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
 AN 1998:281588 BIOSIS
 DN PREV199800281588
 TI The potential of **retro-inverso peptides** as synthetic vaccines.
 AU Van Regenmortel, M. H. V. (1); Guichard, G.; Benkirane, N.; Briand, J.-P.;

Muller, S.; Brown, F.
 CS (1) Inst. Biol. Mol. et Cell., CNRS UPR 9021, 15 rue Rene Descartes,
 F-67084 Strasbourg Cedex France
 SO Brown, F. [Editor]; Haaheim, L. R. [Editor]. Developments in Biological
 Standardization, (1998) Vol. 92, pp. 139-143. Developments in Biological
 Standardization; Modulation of the immune response to vaccine antigens.
 Publisher: S. Karger AG P.O. Box, Allschwilerstrasse 10, CH-4009 Basel,
 Switzerland.
 Meeting Info.: Symposium Bergen, Norway June 18-21, 1996 International
 Association of Biological Standardization
 . ISSN: 0301-5149. ISBN: 3-8055-6640-9.
 DT Book; Conference
 LA English

L4 ANSWER 8 OF 52 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
 AN 1999:16628 BIOSIS
 DN PREV199900016628
 TI A peptide nucleic acid (PNA) is more rapidly internalized in cultured
 neurons when coupled to a retro-inverso delivery peptide. The antisense
 activity depresses the target mRNA and protein in magnocellular oxytocin
 neurons.
 AU Aldrian-Herrada, Gudrun; Desarmenien, Michel G.; Orcel, Helene;
 Boissin-Agasse, Line; Mery, Jean; Brugidou, Jean; Rabie, Alain (1)
 CS (1) CNRS-UPR 9055, Biologie Neurones Endocrines, CCIPE, 141 rue
 Cardonille, 34094 Montpellier Cedex 5 France
 SO Nucleic Acids Research, (Nov. 1, 1998) Vol. 26, No. 21, pp.
 4910-4916.
 ISSN: 0305-1048.
 DT Article
 LA English
 AB A peptide nucleic acid (PNA) antisense for the AUG translation initiation
 region of prepro-oxytocin mRNA was synthesized and coupled to a
retro-inverso peptide that is rapidly taken up
 by cells. This bioconjugate was internalized by cultured cerebral cortex
 neurons within minutes, according to the specific property of the vector
 peptide. The PNA alone also entered the cells, but more slowly. Cell
 viability was unaffected when the PNA concentrations were lower than 10
 muM and incubation times less than for 24 h. Magnocellular neurons from
 the hypothalamic supraoptic nucleus, which produce oxytocin and
 vasopressin, were cultured in chemically defined medium. Both PNA and
 vector peptide-PNA depressed the amounts of the mRNA coding for
 prepro-oxytocin in these neurons. A scrambled PNA had no effect and the
 very cognate prepro-vasopressin mRNA was not affected. The antisense PNA
 also depressed the immunocytochemical signal for prepro-oxytocin in this
 culture in a dose- and time-dependent manner. These results show that PNAs
 driven by the retro-inverso vector peptide are powerful antisense reagents
 for use on cells in culture.

L4 ANSWER 9 OF 52 MEDLINE
 AN 1999097766 MEDLINE
 DN 99097766 PubMed ID: 9881091
 TI A 'retro-inverso' PNA: structural implications for DNA and RNA binding.
 AU Krotz A H; Larsen S; Buchardt O; Eriksson M; Nielsen P E
 CS Center for Biomolecular Recognition, H. C. Orsted Institute, University of
 Copenhagen, Denmark.
 SO BIOORGANIC AND MEDICINAL CHEMISTRY, (1998 Nov) 6 (11) 1983-92.
 Journal code: 9413298. ISSN: 0968-0896.
 CY ENGLAND: United Kingdom
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199902
 ED Entered STN: 19990216
 Last Updated on STN: 19990216

Entered Medline: 19990202

AB **'Retro-inverso' peptide nucleic acid (PNA)**
monomers of thymine (T*: N-(amidomethyl)-N-(N1-thyminyl-acetyl)-beta-alanyl) (and adenine) have been prepared and introduced in PNA oligomers. A homo 'retro-inverso' T*8 PNA was found not to hybridize to a complementary DNA or RNA oligonucleotide, whereas introduction of one retro-inverso thymine unit into the middle of a normal PNA 15-mer resulted in a c.a. 8 degrees C destabilization of the complex of this oligomer with a complementary DNA or RNA oligomer. In an effort to compensate for the structural nucleobase 'phase-shift' caused by the T* monomer by also introducing a beta-alanine monomer it is concluded that the effect of the T* backbone is -7 degrees C when hybridizing to DNA and -4.5 degrees C when hybridizing to RNA. Nonetheless, the T* unit shows good sequence discrimination comparable to that of normal PNA. Molecular dynamics simulations indicate an unfavourable conformation of the backbone amide carbonyl group resulting in reduced interaction with the aqueous medium and an 'electrostatic clash' with the carbonyl of the nucleobase linker. These results show that a simple inversion of an amide bond in the PNA backbone has a dramatic, and hardly predictable, effect on the DNA mimicking properties of the oligomer.

L4 ANSWER 10 OF 52 MEDLINE
AN 1998214886 MEDLINE
DN 98214886 PubMed ID: 9554267
TI The potential of **retro-inverso peptides** as synthetic vaccines.
AU Van Regenmortel M H; Guichard G; Benkirane N; Briand J P; Muller S; Brown F
CS Institut de Biologie Moleculaire et Cellulaire, CNRS UPR, Strasbourg, France.
SO DEVELOPMENTS IN BIOLOGICAL STANDARDIZATION, (1998) 92 139-43.
Journal code: 0427140. ISSN: 0301-5149.
CY Switzerland
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 199806
ED Entered STN: 19980708
Last Updated on STN: 20021218
Entered Medline: 19980625

AB Retro-inverso (RI) peptides, also called all-D-retro peptides, have been shown to mimic the antigenic and immunogenic properties of L-peptides successfully. RI peptides corresponding to the loop 141-159 of the VP1 protein of foot-and-mouth disease virus (FMDV) have been synthesized and used to immunize rabbits and guinea pigs. These peptides induced longer-lasting and higher antibody titres in immunized animals than did the corresponding L-peptides and the antibodies cross-reacted strongly with virus particles and with L-peptides. Antisera raised to RI peptides had in vitro virus neutralization titres equal to or better than those obtained after immunization with classical FMDV antigens and L-peptides. In view of their increased stability, RI peptides may overcome some of the shortcomings of synthetic viral vaccines based on L-peptides.

=> d 11-20 bib ab

L4 ANSWER 11 OF 52 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
AN 1998:414053 BIOSIS
DN PREV199800414053
TI On the potential of **retro-inverso peptides** in vaccine design.
AU Muller, S. (1); Briand, J. P.
CS (1) Inst. Biologie Moleculaire Cellulaire, UPR 9021 CNRS, 15 rue Descartes, F-67000 Strasbourg France

SO Research in Immunology, (Jan., 1998) Vol. 149, No. 1, pp. 55-57.
Meeting Info.: Euroconference on New Trends in Vaccine Research and Development: Adjuvants, Delivery Systems and Antigen Formulations Paris, France February 26-28, 1998
ISSN: 0923-2494.

DT Conference
LA English

L4 ANSWER 12 OF 52 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
AN 1997:465325 BIOSIS
DN PREV199799764528
TI Immunochemical recognition of **retro-inverso**
peptides and their potential as synthetic vaccines.

AU Van Regenmortel, Marc H. V.
CS CNRS UPR 9021, Inst. Biologie Molculaire Cellulaire, F-67084 Strasbourg Cedex France

SO Brown, F. [Editor]; Burton, D. [Editor]; Doherty, P. [Editor]; Mekalanos, J. [Editor]. Vaccines (Cold Spring Harbor), (1997) Vol. 97, pp. 9-15.
Vaccines (Cold Spring Harbor); Molecular approaches to the control of infectious diseases.
Publisher: Cold Spring Harbor Laboratory Press 10 Skyline Drive, Plainview, New York 11803, USA.
Meeting Info.: Fourteenth Annual Meeting on Modern Approaches to the Control of Infectious Diseases Cold Spring Harbor, New York, USA September 9-13, 1996
ISSN: 0899-4056. ISBN: 0-87969-516-1.

DT Book; Conference
LA English

L4 ANSWER 13 OF 52 MEDLINE DUPLICATE 4
AN 1998024168 MEDLINE
DN 98024168 PubMed ID: 9356486
TI A **retro-inverso peptide** corresponding to the
GH loop of foot-and-mouth disease virus elicits high levels of long-lasting protective neutralizing antibodies.

AU Briand J P; Benkirane N; Guichard G; Newman J F; Van Regenmortel M H; Brown F; Muller S
CS Institut de Biologie Molculaire et Cellulaire, Unite Propre de Recherche 9021, Centre National de la Recherche Scientifique, Strasbourg, France.

SO PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (1997 Nov 11) 94 (23) 12545-50.
Journal code: 7505876. ISSN: 0027-8424.

CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 199712
ED Entered STN: 19980109
Last Updated on STN: 19980109
Entered Medline: 19971216

AB Peptides corresponding to the immunodominant loop located at residues 135-158 on capsid protein VP1 of foot-and-mouth disease virus (FMDV) generally elicit high levels of anti-peptide and virus-neutralizing antibodies. In some instances, however, the level of neutralizing antibodies is low or even negligible, even though the level of anti-peptide antibodies is high. We have shown previously that the antigenic activity of peptide 141-159 of VP1 of a variant of serotype A can be mimicked by a retro-inverso (all-D retro or retroenantio) peptide analogue. This retro-inverso analogue induced greater and longer-lasting antibody titers than did the corresponding L-peptide. We now show that a single inoculation of the retro-inverso analogue elicits high levels of neutralizing antibodies that persist longer than those induced against the corresponding L-peptide and confer substantial protection in guinea pigs challenged with the cognate virus. In view of the high stability to

proteases of **retro-inverso peptide** analogues and their enhanced immunogenicity, these results have practical relevance in designing potential peptide vaccines.

L4 ANSWER 14 OF 52 MEDLINE DUPLICATE 5
AN 97249852 MEDLINE
DN 97249852 PubMed ID: 9095678
TI Structural comparison between retro-inverso and parent peptides: molecular basis for the biological activity of a retro-inverso analogue of the immunodominant fragment of VP1 coat protein from foot-and-mouth disease virus.
AU Carver J A; Esposito G; Viglino P; Fogolari F; Guichard G; Briand J P; Van Regenmortel M H; Brown F; Mascagni P
CS Dipartimento di Scienze e Tecnologie Biomediche Universita di Udine, Italy.
SO BIOPOLYMERS, (1997 Apr 15) 41 (5) 569-90.
Journal code: 0372525. ISSN: 0006-3525.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 199705
ED Entered STN: 19970602
Last Updated on STN: 20021218
Entered Medline: 19970521
AB Antibodies induced against intact foot-and-mouth disease Virus (FMDV) particles bind to the retro-inverso analogue of fragment 141-159 of the viral coat protein VP1 of FMDV, variant A, equally well as to the parent peptide. A conformational investigation of this **retro-inverso peptide** was carried out by nmr spectroscopy and restrained molecular modeling in order to identify the structural basis for the antigenic mimicry between these retro-inverso and parent peptides. In 100% trifluoroethanol a well-defined left-handed alpha-helical region exists from residue 150 to residue 159, which is consistently present in all conformational families obtained from restrained modelling. A less-defined left-handed helical region is present in the tract 144-148, which is also consistent for all structures. Conformational flexibility exists about Gly149, which leads to two types of structures, either bent or linear. In the bent structures, a three-residue inverse tight turn is found, which can be classified as an inverse gamma-turn centered at Gly149. The overall structural features of the **retro-inverso peptide** are shown to be similar to those of the parent L-peptide. The two molecules, however, are roughly mirror images because they share inherently chiral secondary structure elements. By comparing these conformational conclusions with the x-ray structure of the Fab complex of a corresponding VP1 antigenic fragment, a rationale is proposed to account for the topological requirements of specific recognition that are implied by the equivalent antigenic activity of the natural and retro-inverso compounds.

L4 ANSWER 15 OF 52 MEDLINE DUPLICATE 6
AN 97295648 MEDLINE
DN 97295648 PubMed ID: 9151257
TI Synthesis and activity of partial retro-inverso analogs of the antimetastatic laminin-derived peptide, YIGSR-NH2.
AU Zhao M; Kleinman H K; Mokotoff M
CS Department of Pharmaceutical Sciences, School of Pharmacy, University of Pittsburgh, Pennsylvania, USA.
NC RR04664-01 (NCRR)
SO JOURNAL OF PEPTIDE RESEARCH, (1997 Mar) 49 (3) 240-53.
Journal code: 9707067. ISSN: 1397-002X.
CY Denmark
DT Journal; Article; (JOURNAL ARTICLE)
LA English

FS Priority Journals
 EM 199707
 ED Entered STN: 19970805
 Last Updated on STN: 19970805
 Entered Medline: 19970724

AB This paper describes the synthesis and biological evaluation of six partial retro-inverso peptidomimetic analogs of YIGSR-NH₂, a synthetic peptide from the beta 1 chain of laminin, which has antimetastatic activity. The intent was to improve the antimetastatic potency of YIGSR-NH₂ by limiting the in vivo enzymatic degradation through the incorporation of fraudulent peptide bonds. We have prepared the following **retro-inverso peptides**, Tyr-Ile-Gly-Ser-gArg-CHO (1), Tyr-gIle-mGly-Ser-Arg-NH₂ (2), Tyr-gIle-mGly-Ser-gArg-CHO (3), gTyr-D-rIle-mGly-Ser-Arg-NH₂ (4), Tyr-Ile-Gly-gSer-D-rArg-CHO (5) and Tyr-gIle-rGly-D-rSer-D-rArg-CHO (6). In vitro assays for B16F10 melanoma cell adhesion showed no significant activity for these six peptides. Peptides 1-3, 5 and 6 were further tested, in vivo, for their ability to inhibit tumor metastases to the lung in mice injected in the tail vein with B16F10 melanoma cells. All five of the **retro-inverso peptides** tested showed statistically significant inhibition of metastasis, but the most active peptides were 5 and 6, which showed 57 and 69% inhibition of metastasis, respectively.

L4 ANSWER 16 OF 52 MEDLINE DUPLICATE 7
 AN 97332614 MEDLINE
 DN 97332614 PubMed ID: 9188848
 TI On the immunogenic properties of **retro-inverso peptides**. Total retro-inversion of T-cell epitopes causes a loss of binding to MHC II molecules.
 AU Herve M; Maillere B; Mourier G; Texier C; Leroy S; Menez A
 CS CEA, Departement d'Ingenierie et d'Etudes des Proteines, CE Saclay, Gif-sur-Yvette, France.
 SO MOLECULAR IMMUNOLOGY, (1997 Feb) 34 (2) 157-63.
 Journal code: 7905289. ISSN: 0161-5890.
 CY ENGLAND: United Kingdom
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199707
 ED Entered STN: 19970721
 Last Updated on STN: 19970721
 Entered Medline: 19970708

AB Retro-inversion is considered an attractive approach for drug and vaccine design since it provides the modified peptides with higher resistance to proteolytic degradation. We therefore investigated in detail the effect of retro-inversion on the immunological properties of synthetic peptides. We have synthesized retro-inverso analogues of MHC II restricted peptides that thus contained the correct orientation of the side chains but an inverse main chain. Retro-inversion made the peptides unable to compete in I E(d) or I A(d) binding tests, demonstrating a very low, if any, capacity to bind to MHC II molecules. These results confirm previous structural data that hydrogen bonds between residues of MHC II molecules and the main chain of antigenic peptides play a major interacting role. In vivo experiments further showed that retro-inversion of a T-cell epitope causes its inability to either sustain in vitro T-cell stimulation or to prime specific T cells. Moreover, the **retro-inverso peptide** was not recognized by antibodies raised against the native peptide and did not elicit antibodies when injected into BALB/c mice. **Retro-inverso peptides** appear to be poor immunogens as a result of their weak capacity to bind to MHC II molecules. As an advantage, they are not expected to trigger undesirable humoral responses such as hypersensitivity or allergic disease. These results also provide a molecular explanation regarding the weak immunogenicity of D-amino acids containing polypeptides.

L4 ANSWER 17 OF 52 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
 AN 1997:175210 BIOSIS
 DN PREV199799466923
 TI Small **retro-inverso peptides** recognize MHC class II HLA, DR(alpha, beta-1*0401).
 AU Howard, Susan C. (1); Zacheis, Michelle L.; Bono, Christine P.; Welply, Joseph K.; Hanson, Gunnar J.; Vuletich, Jennifer L.; Bedell, Louis J.; Summers, Neena L.; Schwartz, Benjamin D.; Woulfe, Susan L.
 CS (1) G.D. Searle Dep. Immunol., St. Louis, MO 63198 USA
 SO Protein and Peptide Letters, (1997) Vol. 4, No. 1, pp. 63-68.
 ISSN: 0929-8665.
 DT Article
 LA English
 AB We have synthesized a series of retro-inverso D-peptide analogues and a peptoid analog that mimic potent seven residue L-peptide ligands for DR(alpha, beta-1*0401). The L-peptide ligands compete against binding of a 13 residue biotinylated ligand, HA307-319 (IC-50 60nM), with competing peptide IC-50 S ranging from 30-200nm. The highest affinity heptamer retro-inverso D-peptide tested gave IC-50 10-mu-M. No binding of the peptoid analog was detected.

L4 ANSWER 18 OF 52 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
 AN 1996:454003 BIOSIS
 DN PREV199699176359
 TI Mimicry of viral epitopes with **retro-inverso peptides** of increased stability.
 AU Benkirane, N. (1); Guichard, G.; Briand, J. P.; Muller, S.; Brown, F.; Van Regenmortel, M. H. V.
 CS (1) Inst. Biol. Mol. Cell., CNRS, 15 rue Descartes, F-67084 Strasbourg France
 SO Brown, F. [Editor]. Developments in Biological Standardization, (1996) Vol. 87, pp. 283-291. Developments in Biological Standardization; New approaches to stabilisation of vaccines potency. Publisher: S. Karger AG P.O. Box, Allschwilerstrasse 10, CH-4009 Basel, Switzerland.
 Meeting Info.: Symposium Geneva, Switzerland May 29-31, 1995
 ISSN: 0301-5149. ISBN: 3-8055-6309-4.
 DT Book; Conference
 LA English

L4 ANSWER 19 OF 52 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
 AN 1997:24961 BIOSIS
 DN PREV199799324164
 TI Synthesis and biological properties of partially modified retro and retro-inverso pseudo peptides of Arg-Gly-Asp (RGD).
 AU Nishikawa, Naoyuki (1); Komazawa, Hiroyuki; Orikasa, Atsushi; Yoshikane, Mitsuo; Yamaguchi, Jiro; Kojima, Masayoshi; Ono, Mitsunori; Itoh, Isamu; Azuma, Ichiro; Fujii, Hideki; Murata, Jun; Saiki, Ikuo
 CS (1) Ashigara Res. Lab., Fuji Photo Film Co. Ltd., Minamiashigara, Kanagawa 250-01 Japan
 SO Bioorganic & Medicinal Chemistry Letters, (1996) Vol. 6, No. 22, pp. 2725-2728.
 ISSN: 0960-894X.
 DT Article
 LA English
 AB Partially modified retro and **retro-inverso peptide** analogs of Arg-Gly-Asp (RGD) were synthesized and examined their inhibitory effects on experimental lung metastasis of murine melanoma and adenosine 5'-diphosphate (ADP) induced platelet aggregation. The analogs showed efficient therapeutic potency for the tumor metastasis but low inhibitory effect on ADP induced platelet aggregation.

L4 ANSWER 20 OF 52 MEDLINE DUPLICATE 8

AN 97020448 MEDLINE
 DN 97020448 PubMed ID: 8866827
 TI Inhibition of angiotensin converting enzyme and potentiation of bradykinin
 by retro-inverso analogues of short peptides and sequences related to
 angiotensin I and bradykinin.
 AU Carmona A K; Juliano L
 CS Department of Biophysics, Escola Paulista de Medicina, Sao Paulo, Brazil.
 SO BIOCHEMICAL PHARMACOLOGY, (1996 Apr 26) 51 (8) 1051-60.
 Journal code: 0101032. ISSN: 0006-2952.
 CY ENGLAND: United Kingdom
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199611
 ED Entered STN: 19961219
 Last Updated on STN: 19961219
 Entered Medline: 19961127
 AB There is pharmacological evidence indicating that, in addition to the
 inhibition of angiotensin converting enzyme (ACE; EC 3.4.15.1), the
 potentiation of bradykinin (BK) responses may also involve the BK receptor
 or some binding site in the structures involved in the contractile
 response to this peptide. Dipeptides such as Val-Trp and some of its
 analogues as well as tripeptide homologues, including total and partial
retro-inverso peptides, were synthesized and
 assayed for their ability to inhibit purified guinea pig plasma ACE and to
 potentiate the action of BK on the isolated ileum of the same species.
 The peptides containing the P2-P1, P1-P'1, and P'1-P'2 inverted amide
 bonds inhibited ACE, were resistant to hydrolysis, and, depending on the
 amino acid composition, some of them potentiated the contractile response
 to BK while others did not. Des-[Arg1]-BK, which has an intrinsic
 activity at concentrations higher than 10^{-5} M, and the very dissimilar
 angiotensin I (AI) analogue [Cys5-Cys10]-angiotensin-I-(5-10)-amide, which
 has no detectable contractile activity, were able to inhibit ACE and
 potentiate BK. In contrast to these peptides, BPP5a and BPP9a from
 Bothrops jararaca venom, and Potentiators B and C from Agkistrodon halys
 blomhoffi venom were more effective as BK potentiators than as ACE
 inhibitors. In conclusion, we have synthesized and assayed compounds that
 preferentially inhibit ACE, e.g. retro-inverso tripeptides, or potentiate
 the response of smooth muscle to BK, e.g. snake venom peptides.

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	24.65	24.86

STN INTERNATIONAL LOGOFF AT 10:28:21 ON 28 JUN 2003

Search Paper 6

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1805JXB

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1		Web Page URLs for STN Seminar Schedule - N. America
NEWS 2		"Ask CAS" for self-help around the clock
NEWS 3	Jun 03	New e-mail delivery for search results now available
NEWS 4	Aug 08	PHARMAMarketLetter(PHARMAML) - new on STN
NEWS 5	Aug 19	Aquatic Toxicity Information Retrieval (AQUIRE) now available on STN
NEWS 6	Aug 26	Sequence searching in REGISTRY enhanced
NEWS 7	Sep 03	JAPIO has been reloaded and enhanced
NEWS 8	Sep 16	Experimental properties added to the REGISTRY file
NEWS 9	Sep 16	CA Section Thesaurus available in CAPLUS and CA
NEWS 10	Oct 01	CASREACT Enriched with Reactions from 1907 to 1985
NEWS 11	Oct 24	BEILSTEIN adds new search fields
NEWS 12	Oct 24	Nutraceuticals International (NUTRACEUT) now available on STN
NEWS 13	Nov 18	DKILIT has been renamed APOLLIT
NEWS 14	Nov 25	More calculated properties added to REGISTRY
NEWS 15	Dec 04	CSA files on STN
NEWS 16	Dec 17	PCTFULL now covers WP/PCT Applications from 1978 to date
NEWS 17	Dec 17	TOXCENTER enhanced with additional content
NEWS 18	Dec 17	Adis Clinical Trials Insight now available on STN
NEWS 19	Jan 29	Simultaneous left and right truncation added to COMPENDEX, ENERGY, INSPEC
NEWS 20	Feb 13	CANCERLIT is no longer being updated
NEWS 21	Feb 24	METADEx enhancements
NEWS 22	Feb 24	PCTGEN now available on STN
NEWS 23	Feb 24	TEMA now available on STN
NEWS 24	Feb 26	NTIS now allows simultaneous left and right truncation
NEWS 25	Feb 26	PCTFULL now contains images
NEWS 26	Mar 04	SDI PACKAGE for monthly delivery of multifile SDI results
NEWS 27	Mar 20	EVENTLINE will be removed from STN
NEWS 28	Mar 24	PATDPAFULL now available on STN
NEWS 29	Mar 24	Additional information for trade-named substances without structures available in REGISTRY
NEWS 30	Apr 11	Display formats in DGENE enhanced
NEWS 31	Apr 14	MEDLINE Reload
NEWS 32	Apr 17	Polymer searching in REGISTRY enhanced
NEWS 33	Jun 13	Indexing from 1947 to 1956 added to records in CA/CAPLUS
NEWS 34	Apr 21	New current-awareness alert (SDI) frequency in WPIDS/WPINDEX/WPIX
NEWS 35	Apr 28	RDISCLOSURE now available on STN
NEWS 36	May 05	Pharmacokinetic information and systematic chemical names added to PHAR
NEWS 37	May 15	MEDLINE file segment of TOXCENTER reloaded
NEWS 38	May 15	Supporter information for ENCOMPAT and ENCOMPLIT updated
NEWS 39	May 16	CHEMREACT will be removed from STN
NEWS 40	May 19	Simultaneous left and right truncation added to WSCA
NEWS 41	May 19	RAPRA enhanced with new search field, simultaneous left and right truncation
NEWS 42	Jun 06	Simultaneous left and right truncation added to CBNB
NEWS 43	Jun 06	PASCAL enhanced with additional data
NEWS 44	Jun 20	2003 edition of the FSTA Thesaurus is now available
NEWS 45	Jun 25	HSDB has been reloaded

NEWS EXPRESS April 4 CURRENT WINDOWS VERSION IS V6.01a, CURRENT
 MACINTOSH VERSION IS V6.0b(ENG) AND V6.0Jb(JP),
 AND CURRENT DISCOVER FILE IS DATED 01 APRIL 2003
 NEWS HOURS STN Operating Hours Plus Help Desk Availability
 NEWS INTER General Internet Information
 NEWS LOGIN Welcome Banner and News Items
 NEWS PHONE Direct Dial and Telecommunication Network Access to STN
 NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
 specific topic.

All use of STN is subject to the provisions of the STN Customer
 agreement. Please note that this agreement limits use to scientific
 research. Use for software development or design or implementation
 of commercial gateways or other similar uses is prohibited and may
 result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 09:40:44 ON 28 JUN 2003

=> file .pub		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 09:41:03 ON 28 JUN 2003

FILE 'BIOSIS' ENTERED AT 09:41:03 ON 28 JUN 2003
 COPYRIGHT (C) 2003 BIOLOGICAL ABSTRACTS INC. (R)

=> s retro inverso
 L1 278 RETRO INVERSO

=> search l1 and peptide
 L2 222 L1 AND PEPTIDE

=> s l2 and py<2001
 L3 187 L2 AND PY<2001

=> duplicate remove l3
 DUPLICATE PREFERENCE IS 'MEDLINE, BIOSIS'
 KEEP DUPLICATES FROM MORE THAN ONE FILE? Y/(N):n
 PROCESSING COMPLETED FOR L3
 L4 115 DUPLICATE REMOVE L3 (72 DUPLICATES REMOVED)

=> d 1-20 bib ab

L4 ANSWER 1 OF 115 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
 AN 2001:230910 BIOSIS
 DN PREV200100230910
 TI Probing SAR of FLRF-NH2 with its N- and C-terminally modified analogs and
retro-inverso peptides.
 AU Kubiak, Teresa M. (1); Larsen, Martha J. (1); Dutton, Fred E. (1);
 Friedman, Alan R. (1)
 CS (1) Animal Health Discovery Research, Pharmacia and Upjohn, Kalamazoo, MI,
 49001 USA
 SO Fields, Gregg B.; Tam, James P.; Barany, George. (2000) pp. 762-763.
 Peptides for the new millennium. print.
 Publisher: Kluwer Academic Publishers 3300 AA, Dordrecht, Netherlands.
 Meeting Info.: 16th American Peptide Symposium Minneapolis, MI, USA June
 26-July 01, 1999

ISBN: 0-7923-6445-7 (cloth).

DT Book; Conference
 LA English
 SL English

L4 ANSWER 2 OF 115 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
 AN 2000:417340 BIOSIS
 DN PREV200000417340
 TI Solid-phase synthesis of partially-modified retro and **retro-inverso** psi(NHCH(CF₃))-**peptides**.
 AU Volonterio, Alessandro (1); Bravo, Pierfrancesco; Moussier, Nathalie; Zanda, Matteo (1)
 CS (1) Dipartimento di Chimica del Politecnico, Via Mancinelli 7, I-20131, Milano Italy
 SO Tetrahedron Letters, (12 August, 2000) Vol. 41, No. 33, pp. 6517-6521. print.
 ISSN: 0040-4039.

DT Article
 LA English
 SL English
 AB The solid-phase synthesis of a novel class of retro and **retro-inverso peptides** featuring a psi(NHCH(CF₃)) surrogate of the classical (NH-CO) retro-**peptide** bond has been accomplished. Wang resin bound alpha-amino esters 2 were engaged in Michael-type N-additions with 3-(E-enoyl)-1,3-oxazolidin-2-one 3, which took place very effectively. Highly chemoselective exocyclic oxazolidinone cleavage, followed by parallel couplings of the resulting polymer bound pseudo-**peptides** 6 with further alpha-amino esters, and final release from the resins 7 delivered a library of nine psi(NHCH(CF₃)) retro and **retro-inverso** pseudo-tripeptides 8 with purity ranging from 75 to > 95%.

L4 ANSWER 3 OF 115 MEDLINE DUPLICATE 1
 AN 2000483575 MEDLINE
 DN 20457139 PubMed ID: 11000007
 TI Design and solution structure of functional **peptide** mimetics of nerve growth factor.
 AU Beglova N; Maliartchouk S; Ekiel I; Zaccaro M C; Saragovi H U; Gehring K
 CS Department of Biochemistry and Montreal Joint Centre for Structural Biology, McGill University, 3655 Promenade Sir William Osler, Montreal, Quebec H3G 1Y6, Canada.
 SO JOURNAL OF MEDICINAL CHEMISTRY, (2000 Sep 21) 43 (19) 3530-40.
 Journal code: 9716531. ISSN: 0022-2623.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 200010
 ED Entered STN: 20001019
 Last Updated on STN: 20001019
 Entered Medline: 20001012

AB The C-D loop in nerve growth factor (NGF) is involved in binding to the NGF receptor, TrkA. It is flexible and adopts several different types conformations in different NGF crystal forms. We have previously shown that a small cyclic **peptide** derived from the C-D loop of NGF binds to the TrkA receptor by mimicking the structure of this loop. To understand structure-function relationships in NGF C-D loop mimetics, we have produced a series of **peptides** predicted to form different types of beta-turns. The **peptides** were tested for their ability to promote cell survival in serum-free medium and to induce TrkA tyrosine phosphorylation. NMR structural studies were used to determine the backbone conformation and the spatial orientation of side chains involved in binding to the TrkA receptor. **Peptides** that form type I or type gammaL-alphaR beta-turns were the most active. The variety of active

loop conformations suggests that the mimetics (and NGF) accommodate the binding site on TrkA by an 'induced fit' mechanism. In agreement with this hypothesis, NMR relaxation measurements detected both fast and slow motion in the **peptides**. We also characterized a **retro-inverso peptide** derived from the NGF C-D loop. This D-amino acid cyclic **peptide** did not adopt a conformation homologous to the NGF C-D loop and was inactive. This may be representative of difficulties in producing structural and functional mimetics by **retro-inverso** schemes.

L4 ANSWER 4 OF 115 MEDLINE
 AN 2000437793 MEDLINE
 DN 20405631 PubMed ID: 10946275
 TI Binding kinetics, structure-activity relationship, and biotransformation of the complement inhibitor compstatin.
 AU Sahu A; Soulika A M; Morikis D; Spruce L; Moore W T; Lambris J D
 CS Protein Chemistry Laboratory, Department of Pathology and Laboratory Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA.
 NC AI 30040 (NIAID)
 CA 16520 (NCI)
 GM 56698 (NIGMS)
 +
 SO JOURNAL OF IMMUNOLOGY, (2000 Sep 1) 165 (5) 2491-9.
 Journal code: 2985117R. ISSN: 0022-1767.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Abridged Index Medicus Journals; Priority Journals
 EM 200009
 ED Entered STN: 20000928
 Last Updated on STN: 20000928
 Entered Medline: 20000919
 AB We have previously identified a 13-residue cyclic **peptide**, Compstatin, that binds to complement component C3 and inhibits complement activation. Herein, we describe the binding kinetics, structure-activity relationship, and biotransformation of Compstatin. Biomolecular interaction analysis using surface-plasmon resonance showed that Compstatin bound to native C3 and its fragments C3b and C3c, but not C3d. While binding of Compstatin to native C3 was biphasic, binding to C3b and C3c followed the 1:1 Langmuir binding model; the affinities of Compstatin for C3b and C3c were 22- and 74-fold lower, respectively, than that of native C3. Analysis of Compstatin analogs synthesized for structure-function studies indicated that 1) the 11-membered ring between disulfide-linked Cys2-Cys12 constitutes a minimal structure required for optimal activity; 2) **retro-inverso** isomerization results in loss of inhibitory activity; and 3) some residues of the type I beta-turn segment also interact with C3. In vitro studies of Compstatin in human blood indicated that a major pathway of biotransformation was the removal of Ile1, which could be blocked by N-acetylation of the **peptide**. These findings indicate that acetylated Compstatin is stable against enzymatic degradation and that the type I beta-turn segment is not only critical for preservation of the conformational stability, but also involved in intermolecular recognition.

L4 ANSWER 5 OF 115 MEDLINE
 AN 2000273399 MEDLINE
 DN 20273399 PubMed ID: 10815952
 TI Determination of biophysical parameters of polypeptide **retro-inverso** isomers and their analogues by capillary electrophoresis.
 AU Hearn M T; Keah H H; Boysen R I; Messana I; Misiti F; Rossetti D V; Giardina B; Castagnola M
 CS Department of Biochemistry and Molecular Biology, Monash University, Clayton, Australia.. milton.hearn@med.monash.edu.au
 SO ANALYTICAL CHEMISTRY, (2000 May 1) 72 (9) 1964-72.

Journal code: 0370536. ISSN: 0003-2700.

CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 200006
ED Entered STN: 20000629

Last Updated on STN: 20000629

Entered Medline: 20000621

AB The relationship between the electrophoretic mobility, microobs, Stokes radius, *rs*, ionization state, and solution conformation of the all L-alpha-polypeptide, 1, the corresponding retro-all D-alpha-polypeptide, 2, and several truncated analogues, 3-5, has been investigated under low pH buffer conditions by high-performance capillary zonal electrophoresis (HPCZE) with coated capillaries. The results confirm that, under these conditions, the all L-alpha-polypeptide, 1, and its **retro-inverso** isomer, 2, exhibit nonidentical electrophoretic mobilities and thus different Stokes radii. At higher pH values, i.e., pH 5.0, the electrophoretic behavior of this **retro-inverso** isomer pair, however, converges. These results indicate that variations in the dipole characteristics of the polypeptide main chain and subtle differences introduced by the spatial constraints of the L-alpha-Pro-->D-alpha-Pro residue replacement lead to differences in the Stokes radii and electrophoretic mobilities of these polypeptides. Since the observed electrophoretic mobilities, microobs, reflect the mean of the mobilities of each charge species participating according to their Stokes radius or their intrinsic charge and mole fraction abundances, the results confirm that polypeptide **retro-inverso** isomers with unmodified amino and carboxy termini are resolvable. This outcome was achieved despite their notional topographical and conformational similarities as assessed from high-field proton nuclear magnetic resonance (1H NMR) spectroscopy and circular dichroism (CD) spectroscopy.

L4 ANSWER 6 OF 115 MEDLINE DUPLICATE 2

AN 2000139754 MEDLINE

DN 20139754 PubMed ID: 10673395

TI Structure-function analysis of the 7B2 CT **peptide**.

AU Apletalina E V; Juliano M A; Juliano L; Lindberg I

CS Department of Biochemistry, Louisiana State University Health Sciences Center, New Orleans, Louisiana, 70112, USA.

NC K02 DA00204 (NIDA)

SO BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, (2000 Jan 27)
267 (3) 940-2.

Journal code: 0372516. ISSN: 0006-291X.

CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 200003
ED Entered STN: 20000320

Last Updated on STN: 20021026

Entered Medline: 20000309

AB Prohormone convertases play important roles in the proteolytic conversion of many protein precursors. The neuroendocrine protein 7B2 and its 31-residue carboxyl-terminal (CT) **peptide** potently and specifically inhibit prohormone convertase 2 (PC2). We have analyzed the residues contributing to inhibition using N-terminal truncation and alanine scanning. Removal of more than 3 residues from the amino-terminal end of CT1-18 resulted in a more than 190-fold drop in inhibitory activity, showing that most of the residues between 3 and 18 are required for inhibition. In agreement, an Ala scan indicated that only 4 residues could be replaced with Ala without losing mid-nanomolar inhibitory potency; in particular, Gln7, Gln9, and Asp12 could be Ala-substituted to yield **peptides** with a similar inhibitory potency to the starting

peptide. The all-d-**retro-inverso**, all-l-**inverso**, and all-d analogues of CT **peptide** were completely inactive, indicating that amino acid side chains and the CT **peptide** main chain interact with PC2. CT **peptide** inhibition could not be competitively blocked by preincubation with truncated CT **peptide** forms, supporting an absolute requirement for the Lys-Lys pair in initial binding of the CT **peptide** to the active site.

Copyright 2000 Academic Press.

L4 ANSWER 7 OF 115 MEDLINE DUPLICATE 3
 AN 2001132004 MEDLINE
 DN 20565873 PubMed ID: 11113331
 TI Prosaptide exacerbates ischemia-induced behavioral deficits in vivo; an effect that does not involve mitogen-activated protein kinase activation.
 AU Lapchak P A; Araujo D M; Shackelford D A; Zivin J A
 CS University of California San Diego, Department of Neuroscience, MTF 316, 9500 Gilman Drive, La Jolla CA 92093-0624, USA.. plapchak@ucsd.edu
 NC NS23814 (NINDS)
 NS28121 (NINDS)
 SO NEUROSCIENCE, (2000) 101 (4) 811-4.
 Journal code: 7605074. ISSN: 0306-4522.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 200103
 ED Entered STN: 20010404
 Last Updated on STN: 20010404
 Entered Medline: 20010301
 AB Prosaposin is a 517 amino acid membrane component and secreted protein(5,7,9) that is proteolytically cleaved to generate the four small glycoproteins; saposins A, B, C and D.(9,13,19) Prosaposin's ability to promote neurite outgrowth(31) and to protect neurons from programmed cell death(28) in vitro, as well as to rescue neurons from ischemia and other damage in vivo(11,12,15,25) implied that prosaposin was neurotrophic/neuroprotectant.(1,7,24,31) The neurotrophic sequence of prosaposin was isolated to smaller **peptide** fragments termed prosaptides(15,31) within the amino terminal portion of saposin C.(1,6,8,10,17,20,21,28) The proposed use of synthetic prosaptides as peripherally administered neuroprotective and/or neurotrophic therapeutic agents has stemmed from their ability to cross the blood-brain barrier,(27) as well as their reported neurotrophic activity in vitro.(15,23,31) Few studies, however, have attempted to characterize these **peptides**, presumably due to their reported instability following peripheral administration.(27) With the recent design of a stable 11-mer **retro-inverso** prosaptide,(15,31) it has become feasible to investigate the pharmacological effects of a stable version of these **peptides** in the validated rabbit spinal cord ischemia model that has been used extensively in the development of therapeutics to treat ischemic stroke.(4,14,16,18) Our results show not only that prosaptide was not neurotrophic/neuroprotectant in vivo, but rather it worsened ischemia-induced behavioral deficits.

L4 ANSWER 8 OF 115 MEDLINE DUPLICATE 4
 AN 2000493919 MEDLINE
 DN 20344114 PubMed ID: 10888201
 TI Synthesis and anti-aggregatory activity of linear **retro-inverso** RGD **peptides**.
 AU Dal Pozzo A; Fagnoni M; Bergonzi R; Vanini L; de Castiglione R; Aglio C; Colli S
 CS G. Ronzoni Institute of Chemical and Biochemical Research, Milan, Italy.. dalpozzo@ronzoni.it
 SO JOURNAL OF PEPTIDE RESEARCH, (2000 Jun) 55 (6) 447-54.

Journal code: 9707067. ISSN: 1397-002X.

CY Denmark
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 200010
ED Entered STN: 20001027
Last Updated on STN: 20001027
Entered Medline: 20001019

AB Six **retro-inverso** tri- and tetrapeptide analogues of RGD were prepared and their anti-aggregatory activity was determined by platelet aggregation tests in comparison with the corresponding parent **peptides**. An efficient method for the introduction of a malonyl-aspartic residue into a **peptide** chain is described for the first time. A 2-3-fold decrease in potency or total loss of bioactivity was observed with the new **peptides**; structure-activity relationships are discussed.

L4 ANSWER 9 OF 115 MEDLINE DUPLICATE 5

AN 2001021901 MEDLINE

DN 20449182 PubMed ID: 10991978

TI **Retro-inverso** prosaptide **peptides** retain bioactivity, are stable In vivo, and are blood-brain barrier permeable.

AU Taylor E M; Otero D A; Banks W A; O'Brien J S

CS Department of Neurosciences, University of California, San Diego, La Jolla, California, USA.

SO JOURNAL OF PHARMACOLOGY AND EXPERIMENTAL THERAPEUTICS, (2000 Oct) 295 (1) 190-4.

Journal code: 0376362. ISSN: 0022-3565.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 200011

ED Entered STN: 20010322

Last Updated on STN: 20010322

Entered Medline: 20001103

AB Prosaptide (trademark of Myelos Corporation, San Diego, CA) **peptides** are based on the 14-amino-acid neurotrophic sequence of human prosaposin and, like the parent protein, have potent neurotrophic and neuroprotective properties. We previously examined the in vivo stability of a series of bioactive Prosaptide **peptides** and designed **peptides** with increased enzymatic stability in the central and peripheral nervous systems. In this article, we examined the stability, biological activity, and permeability of the blood-brain barrier to **retro-inverso** Prosaptide peptidomimetics. Retro-inversion both reverses the primary sequence and replaces L-amino acids with D-amino acids. We examined the bioactivity of five peptidomimetics, Prosaptides D1-D5. Prosaptide D1, a **peptide** containing all D-amino acids with the primary sequence intact, was inactive. However, four **retro-inverso** peptidomimetics, Prosaptides D2-D5 retained bioactivity in neurite outgrowth and [(35)S]GTPgammaS binding assays. We focused on Prosaptide D4 as a prototypical **retro-inverso** Prosaptide peptidomimetic for further study. (125)I-Prosaptide D4 remained intact in brain or serum for 60 min after i.v. administration and was transported across the blood-brain barrier with a unidirectional influx constant of 2.5×10^{-4} ml. g⁻¹. min⁻¹. We conclude that **retro-inverso** Prosaptide peptidomimetics are excellent candidates for development as therapeutics for central nervous system neurodegeneration.

L4 ANSWER 10 OF 115 MEDLINE

DUPLICATE 6

AN 2000092439 MEDLINE

DN 20092439 PubMed ID: 10628817

TI A **retro-inverso** miniantibody with anti-HIV activity.
 AU Levi M; Hinkula J; Wahren B
 CS Department of Virology, Swedish Institute for Infectious Disease Control
 and Karolinska Institute, Solna.. michael.levi@smi.ki.se
 SO AIDS RESEARCH AND HUMAN RETROVIRUSES, (2000 Jan 1) 16 (1) 59-65.
 Journal code: 8709376. ISSN: 0889-2229.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals; AIDS
 EM 200002
 ED Entered STN: 20000229
 Last Updated on STN: 20000229
 Entered Medline: 20000215
 AB An HIV-1-specific miniantibody, a **peptide** representing the third
 heavy chain complementarity-determining region (CDR) of an HIV-specific
 mouse antibody, was characterized and modified with unnatural D-isomeric
 amino acids. The CDR **peptide** and its parent antibody bound to a
 similar epitope, located in the V3 region of HIV-1 gp120. A shortened CDR
 sequence was modified with D-amino acids to create an all-D-amino acid
retro-inverso (RI) **peptide** with a reversed
 sequence order. The RI CDR was less susceptible to proteolytic
 degradation than its L-counterpart and had a higher affinity for HIV-1
peptides. The miniantibody and its parent antibody showed
 neutralization of both primary and laboratory strains of HIV-1. In
 accordance with the binding studies, the RI CDR showed a stronger
 HIV-inhibiting capacity than its L-counterpart. We conclude that the
 anti-HIV **retro-inverso** CDR identified in this study
 has the potential to become a future anti-HIV drug. It has a
 virus-neutralizing capacity in vitro and appears to be stable. Future
 research should focus on characterizing its antiviral activity in vivo.

L4 ANSWER 11 OF 115 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
 AN 2001:89836 BIOSIS
 DN PREV200100089836
 TI Dopaminergic neurons rescued from death by Prosaptide™ D5 in MPTP-treated
 mice.
 AU Liu, J. (1); O'Brien, J. S.
 CS (1) UCSD Sch Med, La Jolla, CA USA
 SO Society for Neuroscience Abstracts, (2000) Vol. 26, No. 1-2, pp. Abstract
 No.-381.24. print.
 Meeting Info.: 30th Annual Meeting of the Society of Neuroscience New
 Orleans, LA, USA November 04-09, 2000 Society for Neuroscience
 . ISSN: 0190-5295.
 DT Conference
 LA English
 SL English
 AB Prosaptide™ D5 is a **retro-inverso** 11-mer
peptide derived from the neurotrophic sequence of Prosaposin in
 the saposin C domain. D5 is stable to protease breakdown and crosses the
 blood-brain barrier intact. We presented in vitro data earlier (639.2,
 29th Neuroscience Meeting, 1999, Miami) that D5 rescued DA neurons from
 MPP+-mediated toxicity; D5 supported cell survival and potentiated neurite
 sprouting in primary DA cultures. In this abstract, D5 rescued DA neurons
 from death in vivo in MPTP-treated C57BL/6 mice. After 24 h treatment with
 MPTP (40 mg/kg, i.p.), D5 was injected i.p. 3 times per week for 2 weeks.
 MPTP treatment decreased the number of DA neurons in the substantia nigra
 (SN) to 30% of controls (P<0.0001 vs controls). The effect of D5 treatment
 was dose-dependent in rescuing DA neurons from death; a dose of 50mg/kg
 increased the number of surviving DA neurons to 63% of controls; a high
 dose of 200mg/kg rescued DA neurons to 93% of control values (P>0.05 vs
 controls). A scrambled **peptide** was ineffective at 200mg/kg.
 Prosaptide™ D5 appears to be a useful agent in the rescue of DA neurons
 and may have therapeutic potential for the therapy of Parkinson's disease.

(Work supported in part by a grant from Myelos Neurosciences to JSO.)

L4 ANSWER 12 OF 115 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
AN 2001:78061 BIOSIS
DN PREV200100078061
TI A stable prosaposin **retro-inverso peptide**
exacerbates ischemia-induced behavioral deficits in rabbits: comparison
with the neuroprotective neurosteroid dehydroepiandrosterone sulfate.
AU Chapman, D. F. (1); Araujo, D. M.; Zivin, J. A.; Lapchak, P. A.
CS (1) UCSD, La Jolla, CA USA
SO Society for Neuroscience Abstracts, (2000) Vol. 26, No. 1-2, pp. Abstract
No.-287.5. print.
Meeting Info.: 30th Annual Meeting of the Society of Neuroscience New
Orleans, LA, USA November 04-09, 2000 Society for Neuroscience
. ISSN: 0190-5295.
DT Conference
LA English
SL English
AB Evidence suggests that prosaposin and neurosteroids can rescue neurons
from ischemic damage and excitotoxicity. Because of their potential
neuroprotective properties, both RIP and DHEAS may be useful in treating
ischemic stroke. We examined the behavioral effects of a stable 11-mer RIP
also known as Prosaptide (all D amino acids: LLEETANNDDL) and DHEAS in
rabbits exposed to reversible spinal cord ischemia produced by temporary
occlusion of the infrarenal aorta; RIP (1 mg/kg) or DHEAS (50 mg/kg) were
administered IV 5 minutes following various durations of aortic occlusion
ranging from 15 to 60 min, which allows for the calculation of the
duration (min) associated with a 50% probability of permanent paraplegia
(P50) for each experimental group. A drug was considered to be
neuroprotective only if it prolonged the P50 compared to the
vehicle-treated control group, which was approximately 25-28 min.
Treatment with RIP significantly ($p<0.05$) decreased the P50 to 20 min (20%
reduction), whereas DHEAS significantly ($p<0.05$) prolonged the P50 to 38
min (35% increase). The prominent neuroprotective effects that were
observed with DHEAS included increased mobility, tactile sensation and
hind limb use. In contrast, RIP exacerbated ischemia-induced behavioral
deficits and increased paraplegia. Overall, our study shows that although
neurotrophic-like properties have been documented for both RIP and DHEAS,
only the latter promotes recovery of spinal cord neuron function following
ischemia, suggesting that it may have therapeutic benefits for the
treatment of ischemic stroke.

L4 ANSWER 13 OF 115 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
AN 2000:292029 BIOSIS
DN PREV200000292029
TI Modulators of beta-amyloid **peptide** aggregation comprising
D-amino acids.
AU Findeis, Mark A. (1); Gefter, Malcolm L.; Musso, Gary; Signer, Ethan R.;
Wakefield, James; Molineaux, Susan; Chin, Joseph; Lee, Jung-J; Kelley,
Michael; Komar-Panicucci, Sonj; Arico-Muendel, Christopher C.; Phillips,
Kathryn; Hayward, Neil J.
CS (1) North Grafton, MA USA
ASSIGNEE: Praecis Pharmaceuticals, Inc., Cambridge, MA, USA
PI US 5985242 November 16, 1999
SO Official Gazette of the United States Patent and Trademark Office Patents,
(Nov. 16, 1999) Vol. 1228, No. 3, pp. No pagination. e-file.
ISSN: 0098-1133.
DT Patent
LA English
AB Compounds that modulate natural beta amyloid **peptide** aggregation
are provided. The modulators of the invention comprise a **peptide**
, preferably based on a beta amyloid **peptide**, that is comprised
entirely of D-amino acids. Preferably, the **peptide** comprises 3-5
D-amino acid residues and includes at least two D-amino acid residues

independently selected from the group consisting of D-leucine, D-phenylalanine and D-valine. In a particularly preferred embodiment, the **peptide** is a **retro-inverso** isomer of a beta amyloid **peptide**, preferably a **retro-inverso** isomer of Abeta17-21. In certain embodiments, the **peptide** is modified at the amino-terminus, the carboxy-terminus, or both. Preferred amino-terminal modifying groups include cyclic, heterocyclic, polycyclic and branched alkyl groups. Preferred carboxy-terminal modifying groups include an amide group, an alkyl amide group, an aryl amide group or a hydroxy group. Pharmaceutical compositions comprising the compounds of the invention, and diagnostic and treatment methods for amyloidogenic diseases using the compounds of the invention, are also disclosed.

L4 ANSWER 14 OF 115 MEDLINE DUPLICATE 7
 AN 1999121111 MEDLINE
 DN 99121111 PubMed ID: 9920919
 TI Solution structure of a **retro-inverso peptide**
 analogue mimicking the foot-and-mouth disease virus major antigenic site.
 Structural basis for its antigenic cross-reactivity with the parent
peptide.
 AU Petit M C; Benkirane N; Guichard G; Du A P; Marraud M; Cung M T; Briand J
 P; Muller S
 CS Laboratoire de Chimie-Physique Macromoléculaire, UMR 7568 CNRS,
 ENSIC-INPL, 54000 Nancy, France.
 SO JOURNAL OF BIOLOGICAL CHEMISTRY, (1999 Feb 5) 274 (6) 3686-92.
 Journal code: 2985121R. ISSN: 0021-9258.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 OS PDB-1BCV; PDB-1BFW
 EM 199902
 ED Entered STN: 19990316
 Last Updated on STN: 20000303
 Entered Medline: 19990226
 AB The antigenic activity of a 19-mer **peptide** corresponding to the
 major antigenic region of foot-and-mouth disease virus and its
 retro-enantiomeric analogue was found to be completely abolished when they
 were tested in a biosensor system in trifluoroethanol. This suggests that
 the folding pattern, which is alpha-helix in trifluoroethanol (confirmed
 by CD measurement), does not correspond to the biologically relevant
 conformation(s) recognized by antibodies. The NMR structures of both
peptides were thus determined in aqueous solution. These studies
 showed that the two **peptides** exhibit similar folding features,
 particularly in their C termini. This may explain in part the
 cross-reactive properties of the two **peptides** in aqueous
 solution. However, the **retro-inverso** analogue appears
 to be more rigid than the parent **peptide** and contains five
 atypical beta-turns. This feature may explain why **retro-**
inverso foot-and-mouth disease virus **peptides** are often
 better recognized than the parent **peptide** by anti-virion
 antibodies.

L4 ANSWER 15 OF 115 MEDLINE DUPLICATE 8
 AN 1999388009 MEDLINE
 DN 99388009 PubMed ID: 10458771
 TI Inhibition of experimental autoimmune encephalomyelitis in SJL mice by
 oral administration of **retro-inverso** derivative of
 encephalitogenic epitope P87-99.
 AU Marino M; Ippolito A; Fassina G
 CS Biopharmaceuticals, TECNOGEN S.C.p.A., Science Park, Piana di Monte Verna,
 Italy.
 SO EUROPEAN JOURNAL OF IMMUNOLOGY, (1999 Aug) 29 (8) 2560-6.
 Journal code: 1273201. ISSN: 0014-2980.

CY GERMANY: Germany, Federal Republic of
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 199909
ED Entered STN: 19990925

Last Updated on STN: 20000303

Entered Medline: 19990914

AB **Retro-inverso** modification of **peptides**

preserves parent **peptide** overall topology and provides at the same time stability to proteolysis, leading to derivatives with prolonged half-life in vitro and in vivo. In this study the encephalitogenic epitope P87 - 99 of myelin basic protein has been prepared in the **retro-inverso** form to examine its biological activity in a murine model of multiple sclerosis. Experiments of in vivo T cell tolerance induction in SJL mice revealed that the **retro-inverso peptide** was able to induce a selective T cell hyporesponsiveness, as measured by a reduction in the proliferative response of lymphnode T cells after antigen challenge. Oral administration of **retro-inverso peptide** decreased the disease severity significantly and delayed considerably the disease onset in treated mice. Enhancement of resistance to proteolysis by **retro-inverso** modification of encephalitogenic epitopes may increase the therapeutic value of oral tolerance induction in the treatment of multiple sclerosis and other Th1-associated inflammatory disorders.

L4 ANSWER 16 OF 115 MEDLINE

DUPLICATE 9

AN 1999447490 MEDLINE

DN 99447490 PubMed ID: 10516644

TI Novel strategies for the design of receptor-selective vasopressin analogues: Aib-substitution and **retro-inverso** transformation.

AU Howl J; Prochazka Z; Wheatley M; Slaninova J

CS Molecular Pharmacology Group, School of Health Sciences, University of Wolverhampton, Wolverhampton WV1 1DJ, UK.

SO BRITISH JOURNAL OF PHARMACOLOGY, (1999 Oct) 128 (3) 647-52.

Journal code: 7502536. ISSN: 0007-1188.

CY ENGLAND: United Kingdom

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199912

ED Entered STN: 20000113

Last Updated on STN: 20000113

Entered Medline: 19991222

AB 1. We determined the pharmacological profile of novel backbone-modified **peptides** designed as protease-resistant, selective analogues of AVP. Binding affinities of **peptides** were determined at both V1A and V2 subtypes of vasopressin receptor (VPR). Biological potencies of selected **peptides** were tested in pressor and antidiuretic bioassays. 2. Substitution of the achiral alpha-aminoisobutyric acid (Aib) at position 4 or 7 of AVP produced **peptides** that selectively bound the V2 VPR. Both [Aib4]AVP (140 IU mg⁻¹) and [Aib7]AVP (36 IU mg⁻¹) are selective antidiuretic agonists with little or no activity in uterotonic and pressor assays. 3. [Aib4] and [Aib7] derivatives of the linear V1A-selective antagonist [PhaaDTyr(Et)2Arg6Tyr(NH2)9]AVP bound selectively and with high affinity (Kd 0.51 and 4.1 nM respectively) to the V1A VPR. Bioassays confirmed that these **peptides** were potent antivasopressor agents (pA2 8.10 and 8.36 respectively). 4. A total **retro-inverso** strategy was used to prepare protease-resistant mimetics of both AVP and linear V1A-selective antagonists. Cyclic **retro-inverso** mimetics of AVP did not bind either V1A or V2 VPRs. In contrast,

rationally designed **retro-inverso** mimetics of linear V1A-selective antagonists selectively bound the V1A VPR. 5. Our findings indicate novel methods to improve the pharmacodynamic and pharmacokinetic parameters of neurohypophysial hormone analogues which could be equally applicable to other **peptide**-receptor systems.

L4 ANSWER 17 OF 115 MEDLINE
 AN 1999258094 MEDLINE
 DN 99258094 PubMed ID: 10326244
 TI Observations on the origin of the non-linear van't Hoff behaviour of polypeptides in hydrophobic environments.
 AU Boysen R I; Wang Y; Keah H H; Hearn M T
 CS Department of Biochemistry and Molecular Biology, Monash University, Clayton, Victoria, Australia.
 SO BIOPHYSICAL CHEMISTRY, (1999 Mar 29) 77 (2-3) 79-97.
 Journal code: 0403171. ISSN: 0301-4622.
 CY Netherlands
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199905
 ED Entered STN: 19990607
 Last Updated on STN: 19990607
 Entered Medline: 19990526
 AB In this paper we describe a general procedure to determine the thermodynamic parameters associated with the interaction of polypeptides or proteins with immobilised lipophilic compounds such as non-polar n-octyl groups. To this end, the binding behaviour of an all L-alpha-polypeptide, 1, and its **retro-inverso**-isomer, 2, has been investigated with an n-octylsilica and water-organic solvent mixture containing different percentages of acetonitrile or methanol over the temperature range of 278-338 K. The results confirm that non-linear van'ts Hoff plots occur with this pair of polypeptide isomers, depending on the solvent composition. These findings are consistent with the changes in the thermodynamic parameters, enthalpy of association, $\Delta H_{\text{assoc},i}$, entropy of association, $\Delta S_{\text{assoc},i}$, and heat capacity, $\Delta C_{p,i}$, all having significant temperature dependencies. Theoretical relationship linking the changes in the $\Delta H_{\text{assoc},i}$, $\Delta S_{\text{assoc},i}$ and $\Delta C_{p,i}$ values of these polypeptide-non-polar ligate systems, as a function of temperature, T, have been validated. Significant differences were observed in the magnitudes of these thermodynamic quantities when acetonitrile or methanol was employed as the organic solvent. The origin of these solvent-dependent effects can be attributed to the hydrogen-bonding propensity of the respective solvent. Involvement of enthalpy-entropy compensation effects associated with the interaction of these polypeptides with the hydrophobic ligates has also been documented. Analysis of empirical extra-thermodynamic relationships associated with molecular structural properties of these polypeptides, such as the slope term, S, derived from the plots of the logarithmic capacity factor, $\log k'_i$, of these polypeptides vs. the volume fraction of the organic solvent, [symbol: see text] as a function of temperature, T, has also revealed similar correlations in terms of the interactive behaviour of polypeptides 1 and 2 under these experimental conditions. These findings provide an extended thermodynamic and extra-thermodynamic framework to examine the solvational, conformational and other equilibrium processes that polypeptides (or proteins) can undergo in the presence of n-alkylsilicas or other classes of immobilised hydrophobic surfaces. The experimental approach utilised in this study with these topologically similar polypeptides thus represents a generic procedure to explore the behaviour of polypeptides or proteins in non-polar environments in terms of their molecular properties and the associated linear free energy relationships that determine their interactive behaviour.

L4 ANSWER 18 OF 115 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

AN 1998:281588 BIOSIS
 DN PREV199800281588
 TI The potential of **retro-inverso peptides** as synthetic vaccines.
 AU Van Regenmortel, M. H. V. (1); Guichard, G.; Benkirane, N.; Briand, J.-P.; Muller, S.; Brown, F.
 CS (1) Inst. Biol. Mol. et Cell., CNRS UPR 9021, 15 rue Rene Descartes, F-67084 Strasbourg Cedex France
 SO Brown, F. [Editor]; Haaheim, L. R. [Editor]. Developments in Biological Standardization, (1998) Vol. 92, pp. 139-143. Developments in Biological Standardization; Modulation of the immune response to vaccine antigens. Publisher: S. Karger AG P.O. Box, Allschwilerstrasse 10, CH-4009 Basel, Switzerland.
 Meeting Info.: Symposium Bergen, Norway June 18-21, 1996 International Association of Biological Standardization
 . ISSN: 0301-5149. ISBN: 3-8055-6640-9.
 DT Book; Conference
 LA English

L4 ANSWER 19 OF 115 MEDLINE DUPLICATE 10
 AN 1998451584 MEDLINE
 DN 98451584 PubMed ID: 9776752
 TI A **peptide** nucleic acid (PNA) is more rapidly internalized in cultured neurons when coupled to a **retro-inverso** delivery **peptide**. The antisense activity depresses the target mRNA and protein in magnocellular oxytocin neurons.
 AU Aldrian-Herrada G; Desarmenien M G; Orcel H; Boissin-Agasse L; Mery J; Brugidou J; Rabie A
 CS CNRS-UPR 9055, Biologie des Neurones Endocrines, CCIPE, 141 rue de la Cardonille, 34094 Montpellier Cedex 5, France.
 SO NUCLEIC ACIDS RESEARCH, (1998 Nov 1) 26 (21) 4910-6.
 Journal code: 0411011. ISSN: 0305-1048.
 CY ENGLAND: United Kingdom
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199812
 ED Entered STN: 19990115
 Last Updated on STN: 19990115
 Entered Medline: 19981223

AB A **peptide** nucleic acid (PNA) antisense for the AUG translation initiation region of prepro-oxytocin mRNA was synthesized and coupled to a **retro-inverso peptide** that is rapidly taken up by cells. This bioconjugate was internalized by cultured cerebral cortex neurons within minutes, according to the specific property of the vector **peptide**. The PNA alone also entered the cells, but more slowly. Cell viability was unaffected when the PNA concentrations were lower than 10 microM and incubation times less than for 24 h. Magnocellular neurons from the hypothalamic supraoptic nucleus, which produce oxytocin and vasopressin, were cultured in chemically defined medium. Both PNA and vector **peptide**-PNA depressed the amounts of the mRNA coding for prepro-oxytocin in these neurons. A scrambled PNA had no effect and the very cognate prepro-vasopressin mRNA was not affected. The antisense PNA also depressed the immunocytochemical signal for prepro-oxytocin in this culture in a dose- and time-dependent manner. These results show that PNAs driven by the **retro-inverso** vector **peptide** are powerful antisense reagents for use on cells in culture.

L4 ANSWER 20 OF 115 MEDLINE
 AN 1999097766 MEDLINE
 DN 99097766 PubMed ID: 9881091
 TI A '**retro-inverso**' PNA: structural implications for DNA and RNA binding.

AU Krotz A H; Larsen S; Buchardt O; Eriksson M; Nielsen P E
 CS Center for Biomolecular Recognition, H. C. Orsted Institute, University of
 Copenhagen, Denmark.
 SO BIOORGANIC AND MEDICINAL CHEMISTRY, (1998 Nov) 6 (11) 1983-92.
 Journal code: 9413298. ISSN: 0968-0896.
 CY ENGLAND: United Kingdom
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199902
 ED Entered STN: 19990216
 Last Updated on STN: 19990216
 Entered Medline: 19990202
 AB 'Retro-inverso' peptide nucleic acid (PNA)
 monomers of thymine (T*: N-(amidomethyl)-N-(N1-thyminy1-acetyl)-beta-
 alanyl) (and adenine) have been prepared and introduced in PNA oligomers.
 A homo 'retro-inverso' T*8 PNA was found not to
 hybridize to a complementary DNA or RNA oligonucleotide, whereas
 introduction of one retro-inverso thymine unit into
 the middle of a normal PNA 15-mer resulted in a c.a. 8 degrees C
 destabilization of the complex of this oligomer with a complementary DNA
 or RNA oligomer. In an effort to compensate for the structural nucleobase
 'phase-shift' caused by the T* monomer by also introducing a beta-alanine
 monomer it is concluded that the effect of the T* backbone is -7 degrees C
 when hybridizing to DNA and -4.5 degrees C when hybridizing to RNA.
 Nonetheless, the T* unit shows good sequence discrimination comparable to
 that of normal PNA. Molecular dynamics simulations indicate an
 unfavourable conformation of the backbone amide carbonyl group resulting
 in reduced interaction with the aqueous medium and an 'electrostatic
 clash' with the carbonyl of the nucleobase linker. These results show
 that a simple inversion of an amide bond in the PNA backbone has a
 dramatic, and hardly predictable, effect on the DNA mimicking properties
 of the oligomer.

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	19.88	20.09

STN INTERNATIONAL LOGOFF AT 09:46:12 ON 28 JUN 2003

Search Page 1

WEST Search History

DATE: Saturday, June 28, 2003

<u>Set Name</u> side by side	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u> result set
<i>DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ</i>			
L3	L2 same eigenvector	1	L3
L2	retro inverso peptide	155	L2
L1	retro inverso same peptide	415	L1

END OF SEARCH HISTORY

WEST

Generate Collection

Print

Search Results - Record(s) 1 through 1 of 1 returned.☐ 1. Document ID: US 20020009756 A1

L3: Entry 1 of 1

File: PGPB

Jan 24, 2002

PGPUB-DOCUMENT-NUMBER: 20020009756

PGPUB-FILING-TYPE: new

DOCUMENT-IDENTIFIER: US 20020009756 A1

TITLE: Algorithmic design of peptides for binding and/or modulation of the functions of receptors and/or other proteins

PUBLICATION-DATE: January 24, 2002

INVENTOR-INFORMATION:

NAME	CITY	STATE	COUNTRY	RULE-47
Mandell, Arnold J.	Asheville	NC	US	
Selz, Karen A.	Asheville	NC	US	
Shlesinger, Michael F.	Rockville	MD	US	

US-CL-CURRENT: 435/7.2; 530/333, 702/19

ABSTRACT:

Methods of designing protein-targeted peptides or peptide analogues whose sequences are derived from the target protein sequences, using target protein sequence, analytically derived templates, and relevant distributions of amino acids for weighted random assignments to those templates. The templates are derived from eigenvectors of the autocovariance matrices of the physicochemically-transformed amino acid sequence of the target proteins; wavelet subsequence templates derived from wavelet transformations of the physicochemically-transformed amino acid sequence of the target proteins; and/or non-overlapping redundant subsequence templates computed from the physicochemically-transformed target protein amino acid sequence. The protein targets include cell receptors; transporters; enzymes; chaperonins; antibodies; surface proteins of infectious agents; and any protein involved in protein-protein interactions. The peptides are designed to bind to and/or otherwise modulate the function of the target protein. Partitioned amino acid distributions for weighted random assignments to the similarly partitioned templates are derived from a variety of physiologically relevant amino acid pools or regions in the target protein sequence relevant to the construction of the templates. Sequential pattern ("mode") matches between candidate peptides and their target proteins are designed such that when examined by maximum entropy, all poles power spectral transformations and/or wavelet transformations, they yield peaks of wavenumbers that differ by $\pm 10\%$ of the larger wavenumber value. Also provided are examples of such mode-matched peptides, as well as methods for their use in elucidating sites on proteins for drug design and testing, detection of disease conditions or contaminants, and as therapeutics for protein function modulation in disease treatment.

L3: Entry 1 of 1

File: PGPB

Jan 24, 2002

DOCUMENT-IDENTIFIER: US 20020009756 A1

TITLE: Algorithmic design of peptides for binding and/or modulation of the functions of receptors and/or other proteins

CLAIMS:

1. A method for synthesizing a peptide based on matching a physicochemical mode of a peptide to the same physicochemical mode of a target polypeptide or protein, followed by synthesizing a retro-inverso peptide version of said peptide comprised of D-amino acids, comprising the steps of: assigning a numerical value of an orderable physicochemical property to each member of a set of peptide constituents, said set of peptide constituents including all the members of the set of naturally-occurring L-amino acids; arranging said peptide constituents in order of said numerical values of said orderable physicochemical property; partitioning said set of peptide constituents into a plurality of peptide constituent groups, whereby each of said peptide constituent groups contains at least one member of said set of peptide constituents, each peptide constituent group encompasses a range of said ordered numerical values, and each member of said set of peptide constituents belongs to only one peptide constituent group; creating a polypeptide physicochemical data series by replacing each amino acid in an amino acid sequence of said target polypeptide or protein with said numerical value of said orderable physicochemical property corresponding to said each amino acid in said amino acid sequence of said target polypeptide or protein; calculating one or more polypeptide eigenvalues and a corresponding polypeptide eigenvector associated with each of said one or more polypeptide eigenvalues by linear decomposition of an autocovariance matrix formed from a sequentially lagged data matrix of said polypeptide physicochemical data series; ordering said one or more polypeptide eigenvalues and said corresponding polypeptide eigenvectors from largest to smallest; selecting one or more of said polypeptide eigenvectors; transforming said one or more of said polypeptide eigenvectors into an eigenvector template; forming a graph of said eigenvector template, wherein said numerical values of said physicochemical property are graphed along the y-axis of said graph and ordered position in said eigenvector template is graphed along the x-axis of said graph; partitioning said graph along said y-axis according to said ranges of said numerical values of said physicochemical property defining said peptide constituent groups, to form a plurality of y-axis ranges; assigning one of said peptide constituents to each position in said peptide by using said graph as a template to create a sequence of a mode-matched peptide, wherein at each ordered position in said eigenvector template along said x-axis of said graph, said one of said peptide constituents assigned to said ordered position has a value of said orderable physicochemical property that is within said y-axis range of said ordered point; determining a sequence of a retro-inverso peptide by inverting said sequence of a mode-matched peptide; and synthesizing said retro-inverso peptide from said sequence, using D-amino acids.

2. A method for synthesizing a peptide based on matching a physicochemical mode of a peptide to the same physicochemical mode of a target polypeptide or protein, followed by synthesizing a retro-inverso version of said peptide comprised of D-amino acids, comprising the steps of: assigning a numerical value of an orderable physicochemical property to each member of a set of peptide constituents, said set of peptide constituents including all the members of the set of naturally-occurring amino acids; arranging said peptide constituents in order of said numerical values of said orderable physicochemical property; partitioning said set of peptide constituents into a plurality of peptide constituent groups, whereby each of said peptide constituent groups contains at least one member of said set of peptide constituents, each peptide constituent group encompasses a range of said ordered numerical values, and each member of said set of peptide constituents belongs to only one peptide constituent group; creating a polypeptide physicochemical data series by replacing each amino acid in an amino acid sequence with said numerical value of said orderable physicochemical property corresponding to said each amino acid in said amino acid sequence; calculating one or more polypeptide eigenvalues and a corresponding polypeptide eigenvector associated with each of said one or more polypeptide eigenvalues by linear decomposition of an autocovariance matrix formed from a sequentially lagged data matrix of said polypeptide physicochemical data series; ordering said one or more polypeptide eigenvalues and said corresponding polypeptide eigenvectors from largest to smallest; selecting one or more of said polypeptide eigenvectors; forming a vector, said vector being a sum of the products of each of said plurality of said polypeptide eigenvectors multiplied by the

corresponding eigenvalue; forming a graph of said vector, wherein said numerical values of said orderable physicochemical property are graphed along the y-axis of said graph, and ordered position in said eigenvector template is graphed along the x-axis of said graph; partitioning said graph along said y-axis according to said range of said numerical values of said orderable physicochemical property defining said peptide constituent groups, to form a plurality of y-axis ranges; and assigning one of said peptide constituents to each position in said peptide by using said graph of said vector as a template, wherein at each ordered position in said eigenvector template along said x-axis of said graph, said one of said peptide constituents assigned to said ordered position has a value of said orderable physicochemical property that is within said y-axis range of said ordered position; determining a sequence of a retro-inverso peptide by inverting said sequence of a mode-matched peptide; and synthesizing said retro-inverso peptide from said sequence, using D-amino acids.

Full	Title	Citation	Front	Review	Classification	Date	Reference	Sequences	Attachments	Claims	NMC	Draw Desc	Image
------	-------	----------	-------	--------	----------------	------	-----------	-----------	-------------	--------	-----	-----------	-------

[Generate Collection](#)[Print](#)

Term	Documents
EIGENVECTOR	1158
EIGENVECTORS	1331
(2 SAME EIGENVECTOR).USPT,PGPB,JPAB,EPAB,DWPI.	1
(L2 SAME EIGENVECTOR).USPT,PGPB,JPAB,EPAB,DWPI.	1

Display Format: [REV, K](#) [Change Format](#)[Previous Page](#)[Next Page](#)